

Numerical Methods Revision

| | | |
|--|--|--|
| (a) | (b) | (c) |
| $x_{n+1} = \frac{e^{x_n}}{x_n + 3}$ <p>Using $x_1 = 1$ find the values of x_2, x_3 and x_4 to 4 decimal places.</p> | $f(x) = \ln(2x) - \frac{3}{x}$ <p>The equation $f(x) = 0$ has a single root α. Show that α lies in the interval $[2, 2.1]$.</p> | $f(x) = x^3 - x - 2 \sin x$ <p>Show that the equation $f(x) = 0$ can be written as the iterative formula</p> $x_{n+1} = \sqrt{1 + \frac{2 \sin x_n}{x_n}}$ |
| (d) | (e) | (f) |
| <p>Show that the equation $x^3 - \sqrt{x} = 3$ has the solution $x = 1.623$ to 3 decimal places.</p> | $x_{n+1} = 2 + 3 \ln x_n$ <p>Using $x_0 = 6$, find x_1, x_2 and x_3. Hence find the root of the equation $2 + 3 \ln x - x = 0$ to 3 decimal places.</p> | <p>The iterative formula</p> $x_{n+1} = 5 + 2 \ln \left(\frac{1}{x_n + 3} \right)$ <p>is used to find an approximate solution to the equation $5 - x + 2 \ln \left(\frac{1}{x+3} \right) = 0$</p> <p>Given $x_1 = 1$, show using a cobweb or staircase diagram how the convergence towards the root takes place.</p> |
| (g) | (h) | |
| <p>Use the Newton-Raphson method with $x_1 = 1$ to find an approximate root to the equation $5 + x - e^x = 0$ to 3 decimal places.</p> | <p>The height of a ball as a function of time can be modelled by the equation</p> $h(t) = 5\sqrt{t} - \frac{1}{2}t^3 \quad t \geq 0$ <p>Taking a first approximation of $t = 2$ s, use the Newton-Raphson method once to find a second approximation for the time when the ball hits the ground. Give your answer to 3 significant figures.</p> | |